

# Instructions for Use

**RUZURGI (rew-ZUR-jee)**  
(amifampridine)

For patients who require a dosage in less than 5 mg increments, have difficulty swallowing tablets, or require a feeding tube, a 1 mg/mL suspension can be prepared.

## Supplies Needed:

Supplies are available at your local pharmacy



Empty 30 mL Bottle



Sterile Water



10 mL Oral Syringe  
with a Catheter Tip

## Procedure for making a 1 mg/mL suspension for patients requiring 10 mg or less for each dose

Do not use any foods or liquids other than sterile water to mix RUZURGI.



Place one (1) 10 mg Ruzurgi (amifampridine) tablet in a 30 mL bottle.

### Step 1




Fill an oral syringe with 10 mL of sterile water and inject the contents into the 30 mL bottle.

### Step 2



Secure the cap onto the bottle and shake well for 30 seconds.

### Step 3



Place three (3) 10 mg Ruzurgi (amifampridine) tablets in a 30 mL bottle.

### Step 1



Fill an oral syringe with 10 mL of sterile water and inject the contents into the 30 mL bottle. This step must be performed for a total of three (3) times to create a volume of 30 mL which is equal to a 30 mg dose.

### Step 2



Secure the cap onto the bottle and shake well for 30 seconds.

### Step 3


Prepare fresh suspensions daily. Refrigerate the solution between doses, shaking well before drawing up each dose.

## Procedure for administering RUZURGI by mouth



Remove the bottle cap and use an oral syringe with a catheter tip to measure the prescribed dose.

### Step 4



Administer by mouth. For patients requiring more than 10 mg for each dose, repeat steps 4 and 5 until the prescribed dose is given.

### Step 5

## Procedure for administering RUZURGI through a feeding tube

**Use only an oral syringe with a catheter tip to give RUZURGI through the feeding tube.** Talk to your doctor about the size catheter tipped syringe you should use. Do not use any foods or liquids other than sterile water to mix RUZURGI.



Remove the bottle cap and use an oral syringe with a catheter tip to measure the prescribed dose.

### Step 4



Inject the medicine using the oral syringe with a catheter tip into the feeding tube right away. For patients requiring more than 10 mg for each dose, repeat steps 4 and 5 until the prescribed dose is given.

### Step 5



To flush the feeding tube, refill the syringe with 10 mL of sterile water.

### Step 6



Shake the syringe, insert the catheter tip into the feeding tube to flush any remaining medicine from the feeding tube into the stomach.

### Step 7

## How should I store RUZURGI?

- Store RUZURGI tablets in the container from the pharmacy at room temperature between 68°F to 77°F (20°C to 25°C) for up to 3 months.
- Refrigerate prepared RUZURGI oral suspension between doses.
- Safely throw away unused RUZURGI oral suspension after 24 hours.
- **Keep RUZURGI and all medicines out of the reach of children.**